RVPublisher
a better way to publish

Kaveh Bazargan

23 April 2015

www.RiverValleyTechnologies.com @kaveh1000

25 years of typesetting

- Convert XML to PDF *automatically*
  - Including complex math
- Why not put system into the cloud
- Let others do the work
- …and the computer do the admin

Typical workflow



Wouldn’t it be nice…



Wouldn’t it be nice…



Wouldn’t it be nice…



rvpublisher.rivervalleytechnologies.com

transistor had a low gain of only 0.24 [7]. Yet many devices, such as amplifiers and repeaters, work by feeding the output of one transistor into the input of another transistor.

Hide changes | Add Note | Check In

View | Changes | Approve for Publication

B I x^2 x_2 Save Cancel

The key to achieving gains above unity is creating strong photon-photon interactions. To make these interactions, the single-photon transistors demonstrated by the Rempe and Hofferberth groups rely on optically excited atoms in high-excitation states, called Rydberg states, and an effect called Rydberg blockade that occurs in clouds of identical atoms. An atom excited to a Rydberg state has a large electric dipole, which induces shifts in the energy levels of neighboring atoms within a so-called "blockade radius." As a result, these neighboring atoms become opaque or transparent to a subsequent light source, depending on how this source's wavelength is tuned relative to the light that excited the Rydberg atom. This interaction persists within a so-called "blockade radius," within which other Rydberg excitations are prevented from occurring (Fig. 1). The Rydberg blockade effect accordingly mediates a strong interaction between the excitation photon and the source photons.

FIG. 1. A photon transistor uses a light source to modulate a medium's opacity to subsequent photons. To make a medium whose opacity can be switched with a single photon, researchers used highly excited atoms in an atom cloud. When an incoming photon (red) excites an atom to a Rydberg state (red sphere), the excited atom perturbs the atoms within a certain "blockade radius" (purple volume) rendering them either opaque or transparent to a subsequent light source.



The interaction between two Rydberg atoms can be many orders of magnitude larger than



and their collaborators shows that functional single-photon transistors are possible, and illustrates the general power of Rydberg quantum optics for designing prototype photonic devices.

References

[1]
J. D. Pritchard, D. Maxwell, A. Gauguet, K. J. Weatherill, M. P. A. Jones, and C. S. Adams
Cooperative Atom-Light Interaction in a Blockaded Rydberg Ensemble.
Phys. Rev. Lett. 105 193603 2010
DOI: 10.1103/PhysRevLett.105.193603

[2]
Y. O. Dudin and A. Kuzmich
Strongly Interacting Rydberg Excitations of a Cold Atomic Gas.
Science 336 887 2012
DOI: 10.1126/science.1217901

[3]
O. Firstenberg, T. Peyronel, Q.-Y. Liang, A. V. Gorshkov, M. D. Lukin, and V. Vuletic
Attractive Photons in a Quantum Nonlinear Medium.
Nature 502 71 2013
DOI: 10.1038/nature12512

[4]
Daniel Tiarks, Simon Baur, Katharina Schneider, Stephan Dürr, and Gerhard Rempe
Single-Photon Transistor Using a Förster Resonance.
Phys. Rev. Lett. 113 053602 2014
CODEN: PRLTAO
DOI: 10.1103/PhysRevLett.113.053602

[5]
H. Gorniaczyk, C. Tresp, J. Schmidt, H. Fedder, and S. Hofferberth
Single-Photon Transistor Mediated by Interstate Rydberg Interactions.
Phys. Rev. Lett. 113 053601 2014

rvpublisher.rivervalleytechnologies.com

kaveh

Hide changes | Add Note | Check In
View | Changes | Approve for Publication

transistor had a low gain of only 0.24 [7]. Yet many devices, such as amplifiers and repeaters, work by feeding the output of one transistor into the input of another transistor. Such devices require transistors with gains above unity.

The key to achieving gains above unity is creating strong photon-photon interactions. To make these interactions, the single-photon transistors demonstrated by the Rempe and **Hofferberth** groups rely on optically excited atoms in high-excitation states, called Rydberg states, and an effect called *Rydberg* blockade that occurs in clouds of identical atoms. An atom excited to a Rydberg state has a large electric dipole, which induces shifts in the energy levels of neighboring atoms within a so-called "blockade radius." As a result, these neighboring solid state atoms become opaque or transparent to a subsequent light source, depending on how this source's wavelength is tuned relative to the light that excited the Rydberg atom. ~~This~~ That interaction persists within a so-called "blockade radius," within which other Rydberg excitations are prevented from occurring (Fig. 1). The Rydberg blockade effect accordingly mediates a strong interaction between the excitation photon and the source photons.

Added bold

FIG. 1. A photon transistor uses a light source to modulate a medium's opacity to subsequent photons. To make a medium whose opacity can be switched with a single photon, researchers used highly excited atoms in an atom cloud. When an incoming photon (red) excites an atom to a Rydberg state (red sphere), the excited atom perturbs the atoms within a certain "blockade radius" (purple volume) rendering them either opaque or transparent to a subsequent light source.



rvpublisher.rivervalleytechnologies.com

Hide changes | Add Note | Check In
View | Changes | Approve for Publication

transistor had a low gain of only 0.24 [7]. Yet many devices, such as amplifiers and repeaters, work by feeding the output of one transistor into the input of another transistor. Such devices require transistors with gains above unity.

The key to achieving gains above unity is creating strong photon-photon interactions. To make these interactions, the single-photon transistors demonstrated by the Rempe and **Hofferberth** groups rely on optically excited atoms in high-excitation states, called Rydberg states, and an effect called *Rydberg* blockade that occurs in clouds of identical atoms. An atom excited to a Rydberg state ~~has a large~~ space-indendent electric dipole, which induces shifts in the energy levels of neighboring atoms within a so-called "blockade radius." As a result, these neighboring solid state atoms become opaque or transparent to a subsequent light source, depending on how this source's wavelength is tuned relative to the light that excited the Rydberg atom. ~~This~~ That interaction persists within a so-called "blockade radius," within which other Rydberg excitations are prevented from occurring (Fig. 1). The Rydberg blockade effect accordingly mediates a strong interaction between the excitation photon and the source photons. Some more text added here.

Replace Figure | Remove Figure

FIG. 1. A photon transistor uses a light source to modulate a medium's opacity to subsequent photons. To make a medium whose opacity can be switched with a single photon, researchers used highly excited atoms in an atom cloud. When an incoming photon (red) excites an atom to a Rydberg state (red sphere), the excited atom perturbs the atoms within a certain "blockade radius" (purple volume) rendering them either opaque or transparent to a subsequent light source.



rvpublisher.rivervalleytechnologies.com

achieved all-optical switching in an atomic system with a single gate photon, but this transistor had a low gain of only 0.24 [7]. Yet many devices, such as amplifiers and repeaters, work by feeding the output of one transistor into the input of another transistor. Such devices require transistors with gains above unity.

Hide changes | Add Note | Check In

View | Changes | Approve for Publication

The key to achieving gains above unity is creating strong photon-photon interactions. To make these interactions, the single-photon transistors demonstrated by the Rempe and **Hofferberth** groups rely on optically excited atoms in high-excitation states, called Rydberg states, and an effect called *Rydberg* blockade that occurs in clouds of identical atoms. An atom excited to a Rydberg state ~~has a large~~ space-indendent electric dipole, which induces shifts in the energy levels of neigh ockade radius." As a result, these neighboring solid sta rent to a subsequent light source, depending on how th tive to the light that excited the Rydberg atom. ~~This~~ T alled "blockade radius," within which other Rydbe ccurring (Fig. 1). The Rydberg blockade effect accordin tween the excitation photon and the source photons. S

Action

Changes made by sw 1

Added text

Accept | Reject | Cancel

FIG. 1. A photon transistor uses a light source to modulate a medium's opacity to subsequent photons. To make a medium whose opacity can be switched with a single photon, researchers used highly excited atoms in an atom cloud. When an incoming photon (red) excites an atom to a Rydberg state (red sphere), the excited atom perturbs the atoms within a certain "blockade radius" (purple volume) rendering them either opaque or transparent to a subsequent light source.



rivervalleytechnologies.com/view-stf-article?ai=12&sti=39

rvpublisher.rivervalleytechnologies.com

achieved all-optical switching in an atomic system with a single gate photon, but this transistor had a low gain of only 0.24 [7]. Yet many devices, such as amplifiers and repeaters, work by feeding the output of one transistor into the input of another transistor. Such devices require transistors with gains above unity.

The key to achieving gains above unity is creating strong photon-photon interactions. To make these interactions, the single-photon transistors demonstrated by the Rempe and Hofferberth groups rely on optically excited atoms in high-excitation states, called Rydberg states, and an effect called Rydberg blockade that occurs in clouds of identical atoms. An atom excited to a Rydberg state has a large space-indendent electric dipole, which induces shifts in the energy levels of neighboring atoms within a so-called "blockade radius." As a result, these neighboring solid state atoms become opaque or transparent to a subsequent light source, depending on how this source's wavelength is tuned relative to the light that excited the Rydberg atom. This That interaction persists within a so-called "blockade radius," within which other Rydberg excitations are prevented from occurring (Fig. 1). The Rydberg blockade effect accordingly mediates a strong interaction between the excitation photon and the source photons. Some more text added here.

FIG. 1. A photon transistor uses a light source to modulate a medium's opacity to subsequent photons. To make a medium whose opacity can be switched with a single photon, researchers used highly excited atoms in an atom cloud. When an incoming photon (red) excites an atom to a Rydberg state (red sphere), the excited atom perturbs the atoms within a certain "blockade radius" (purple volume) rendering them either opaque or transparent to a subsequent light source.



Hide changes
Add Note
Check In
View
Changes
Approve for Publication
XML
PDF

Physics_7_7 (6).pdf (page 1 of 3)
Search

River Valley Research **7**, 7 (2014)

Review

Statistical flickers in a Bose-Einstein Condensate of Photons Phonons

Cristiano Ciuti
Laboratoire Matériaux et Phénomènes Quantiques,
Université Paris Diderot et CNRS, Paris, France

Published January 21, 2014

Experiments with photons show that the number of particles in a Bose-Einstein condensate can have strong fluctuations.

Subject Areas: **Optics, Statistical Physics**

Summary of:
Statistical flickers in a Bose-Einstein Condensate of Photons Phonons
Journal Info – January 21, 2014

The most familiar type of Bose-Einstein condensate (BEC) is one software writer text added made of identical atoms, which, when cooled to extremely low temperatures, collectively fall into the same single-particle quantum state and form a giant matter wave. But BECs can form from other types of ***bosons*** , including polaritons and photons, and unlike atoms in an isolated trap, these particles can be easily created or destroyed within the medium in which they are produced. Researchers have now revealed unusual statistical effects in a BEC that can exchange particles and energy with such a medium. Julian Schmitt and colleagues at the University of Bonn, Germany, produced a BEC of photons within a bath of photoexcitable dye molecules that acted as a "reservoir" of photons and a heat bath, showing that particle fluctuations in the condensate could be a substantial fraction of the total number of particles [1]. Their experiment, reported in *Physical Review Letters*, is a rare example of a quantum gas in the so-called grand-canonical ensemble of statistical physics. Their work also shows that it served and large fluctuations aren't allowed. Similarly, large fluctuations shouldn't occur in a "microcanonical" ensemble, in which both energy and particle number are conserved.

So far, experimentalists haven't seen these ensemble-dependent statistical effects in BECs. One reason is practical: in the case of an atom BEC, the atomic gas has to be isolated from its environment in order to cool it to low temperatures. These BECs are therefore usually prepared in microcanonical ensemble conditions [2]. In principle, it should be possible to see particle fluctuations in the BECs of exciton-polaritons [3] that can occur in solid-state systems. But both the accurate control of the reservoir and a precise measurement of the fast (picosecond-scale) fluctuations have proven difficult. Finally, theoretical work has questioned if it is possible for a BEC to form under grand-canonical ensemble conditions [4].

The Bonn group shows that, in fact, these conditions

www.RiverValleyTechnologies.com @kaveh1000

[2] For a comprehensive review with historic insights, see V. V. Kocharovsky, V. V. Kocharovsky, M. Holthaus, C. H. R. Ooi, A. Svidzinsky, W. Ketterle, and M. O. Scully, "Fluctuations in Ideal and Interacting Bose-Einstein Condensates: From the Laser Phase Transition Analogy to Squeezed States and Bogoliubov Quasiparticles," *Adv. At. Mol. Opt. Phys.* **53**, 291 (2006).

[3] J. Kasprzak, "Bose-Einstein Condensation of Exciton Polaritons," *Nature* **443**, 409 (2006).

[4] R. M. Ziff, G. E. Uhlenbeck, and M. Kac, "The Ideal Bose-Einstein Gas, Revisited," *Phys. Rep.* **32**, 169 (1977).

[5] J. Klaers, J. Schmitt, F. Vewinger, and M. Weitz, "Bose-Einstein Condensation of Photons in an Optical Microcavity," *Nature* **468**, 545 (2010).

[6] J. Klaers, J. Schmitt, T
"Statistical Physics of
Dye Microcavity," *Phys*

[7] D. Bajoni, P. Senellart,
Lasing in a GaAs Micr
Condensate," *Phys. Rev*

[8] B. Fischer and R. Weill,
come a Condensation Pl
(2012).

[9] P. Kirton and J. Keelin
Condensation," *Phys. R*

[10] I. Carusotto and C. Ciu
Mod. Phys. **85**, 299 (20

Why we use LaTeX

- LaTeX is not just for math
  - Best line breaking, page breaking…
  - Highest quality typography
- Batch process, so automated
- Interactive PDFs direct from XML

Enhanced tooltips



Self-modifying PDF



Reader-customized PDFs




PeerJ
Show data
Metrics
About the author
Reviews
Comments
Transcriptome analyses of Anguillicola crassus from native and novel hosts
Usage since published - updated daily
Social referrals unique visitors
Twitter 26
Google+ 0
Facebook 86
LinkedIn 0
Reddit 0
Slashdot 0
Social networks
Tweet 14
Recommend 42
Share 0
Top referrals unique visitors
dml.cmnh.org 25
From bookmark or typed URL 21
Close
introduction. In the new host in the recipient area, the parasite appears to be more pathogenic. As a reason for these differences, genetically fixed differences in infectivity and development between Taiwanese and European A.crassus have been described and disentangled from plasticity induced by different host environments. To explore whether transcriptional regulation is involved in these lifecycle differences, we have analysed a "common garden", cross infection experiment, using deep-sequencing

Let's ditch attachments



Time's Up!

About your speaker:

Name: ***Kaveh Bazargan***

Company: ***River Valley Technologies***

Tel: *+44 7771 824 111*

Email: ***kaveh@RiverValleyTechnologies.com***

Social Media: ***kaveh1000***